

## **DEPARTMENT OF SOCIAL SERVICES**

DIVISION OF MEDICAL SERVICES 700 GOVERNORS DRIVE PIERRE, SD 57501-22941

**PHONE:** 605-773-3495 | FAX: 605-773-5246

WEB: DSS Medicaid Prior Authorizations | EMAIL: DSSMedicaidpa@state.sd.us

## **EVKEEZA PRIOR AUTHORIZATION REQUEST FORM**

This form **MUST BE** submitted with medical records to support services

Date:					
RECEIPIENT INFORMATION					
Medicaid ID:	Date of Birth:		Sex: M F		
Last Name:		First Name:			
GENERAL INFORMATION					
First Date of Service:		Last Date of Service:			
Primary Diagnosis Code:		HCPC Code:			
Drug Name:		Dose & Frequency:			
Hospitalizations/Treatments/Medications Used in the last 6 months:					
	DOINT OF	CONTACT			
POINT OF CONTACT					
Name and Title:					
Email:	Phone:		Fax:		
Note: The point of contact is the individual completing the PA and would be the contact for questions SD Medicaid may have regarding the PA. The determination notice will be sent to the listed point of contact.					
REFERRING PROVIDER INFORMATION					
Name:					
NPI #:		Taxonomy:			
Phone:		Fax:			
SERVICING PROVIDER INFORMATION					
Name:					
Address:					
NPI #:		Taxonomy:			
Phone:		Fax:			

CRITERIA					
Medical records to support use of product are submitted					
Initial Therapy (check one)	Yes	No			
Therapy is prescribed by or in	। consultation with a cardio	ologist, endocrinologist, or lipid specialist			
Individual has a diagnosis of HoFH defined as <b>one</b> of the following:  • Genetic mutation indicating HoFH (e.g., mutations in low density lipoprotein receptor [LDLR] gene, proprotein convertase subtilisin kexin 9 [PCSK9] gene, apolipoprotein B [apo B] gene, low density lipoprotein receptor adaptor protein 1 [LDLRAP1] gene)  • Treated LDL-C ≥ 300 mg/dL and one of the following  • Tendinous or cutaneous xanthoma prior to age 10 years  • Evidence of HeFH in both parents (e.g., documented history of elevated DLC ≥ 190 mg/dL prior to lipid-lowering therapy)  • Untreated LDL-C ≥ 500 mg/dL  Individual meets <b>one</b> of the following regarding statin therapy  • Individual is currently taking high intensity statin therapy (high intensity statin is defined at atorvastatin ≥40mg or rosuvastatin ≥20mg) and has been compliant with therapy for ≥3 months  • Individual is statin intolerant based on <b>one</b> of the following					
daily starting dos  Development of necrotizing myor  Individual has a contra	se statin associated rhabdo pathy (IMNM) after a trial iindication to statin therap	th at least one started at the lowest myolysis or immune-mediated of one statin by including but not limited to active liver disease, saminases or pregnancy			
months  Individual has had a tri		daily and has been compliant with therapy for ≥3  nse to ezetimibe therapy			
Individual meets <b>one</b> of the folionhibitor therapy  Individual is currently to FDA approved doses are individual has had a tricof ≤50% from baseline  Individual is intolerant  Documentation is prov	lowing regarding proprote aking PCSK9 therapy with and has been compliant with all and inadequate respo of PCSK9 therapy ided that individual is LD	ein convertase subtilisin kexin type 9 (PCSK9)  th evolocumab (Repatha) or alirocumab (Praluent) at vith therapy for ≥3 months.  nse to PCSK9 therapy as indicated by LDL reduction  LR negative			
Therapy will not be used concomitantly with lomitapide (Juxtapid)					
Individual is ≥5 years of age					

Continuation of Therapy (check one)	Yes	No		
Individual continues to meet initia	al criteria			
		ipid lowering therapies including maximum tolerated previously waived due to contraindication or		
Documentation is provided individual has had a reduction in total LDL-C since starting Evkeeza therapy				
PHYSICIAN SIGNATURE – PROVIDER ONLY				
This form must be signed by a provider				
I certify that the information giver product	າ in this form is a true	and accurate medical indication for the required		
Name & Title (Printed):		Specialty:		
Signature:		,		