



# Well-Child Check-up REMINDER



**To the parents of:**

Your child may be due for a well-child check-up. Doctors recommend infants and toddlers have 12 check-ups by age 3. Your child should have a check-up at the following ages:

3-5 Days	4 Months	12 Months	24 Months
1 Month	6 Months	15 Months	30 Months
2 Months	9 Months	18 Months	36 Months

Well-child check-ups help identify health concerns early and keep your child healthy. In addition, your child may be due for a recommended vaccine. Vaccines can help prevent serious health problems including diseases that can be fatal.

Well-child check-ups and vaccines are covered as part of your child's Medicaid/CHIP coverage at no cost to you. Based on the age of your child, the following services may be recommended:

---

You can schedule a check-up by contacting your child's Primary Care Provider (PCP). If you think your child may be behind on his or her vaccines, you should discuss whether any additional vaccines are needed with your child's PCP. Check-ups are also a good time to discuss with your child's PCP any stress, questions, or concerns you have as a parent of a young child.


Dental check-ups are also covered as part of your child's Medicaid/CHIP coverage. Dentists recommend children have a dental check-up around age 1 and every 6 months after that. To find a dentist near you visit <https://www.insurekidsnow.gov/find-a-dentist/index.html>.



Department of Social Services - Division of Medical Services  
700 Governors Drive, Pierre, SD 57501  
Ph: 1.800.597.1603 | Fax: 605.773.5246  
[dss.sd.gov/medicaid/recipients/](https://dss.sd.gov/medicaid/recipients/)

# 0 to 3-year-old WELL-CHILD CHECK-UPS

## WELL VISIT AND IMMUNIZATION ROAD MAP



Vaccine	Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	19-23 months	2-3 years
<b>HepB</b> (Hepatitis B)	1 dose	1 dose				1 dose				
<b>RV</b> (Rotavirus)			1 dose	1 dose	1 dose					
<b>DTap</b> (Tetanus, diphtheria, pertussis)			1 dose	1 dose	1 dose		1 dose			
<b>Hib</b> (Haemophilus influenzae type b)			1 dose	1 dose	1 dose	1 dose				
<b>PCV</b> (Pneumococcal)			1 dose	1 dose	1 dose	1 dose				
<b>IPV</b> (Polio)			1 dose	1 dose		1 dose				
<b>Flu</b> (Influenza)						2 doses in first year, then 1 dose annually				
<b>MMR</b> (Measles, Mumps, Rubella)						1 dose				
<b>Varicella</b> (Chickenpox)						1 dose				
<b>HepA</b> (Hepatitis A)						2 doses				

